



### Pupil Gauge (in mm)



### GLASGOW COMA SCALE

INFANT		Eye Opening	CHILD/ ADULT
4	Spontaneously	Spontaneously	4
3	To speech	To speech	3
2	To pain	To pain	2
1	No response	No response	1
Best Verbal Response			
5	Coos, babbles	Oriented	5
4	Irritable cries	Confused	4
3	Cries to pain	Inapprop. words	3
2	Moans, grunts	Incomprehensible	2
1	No response	No response	1
Best Motor Response			
6	Spontaneous	Obeys commands	6
5	Localizes pain	Localizes pain	5
4	Withdraws from pain	Withdraws from pain	4
3	Flexion (decorticate)	Flexion (decorticate)	3
2	Extension (decerebrate)	Extension (decerebrate)	2
1	No response	No response	1
_____ = TOTAL*		*TOTAL = _____	

**\*GCS ≤ 8? - Intubate!**

### RAPID TRIAGE

(for multiple patient scenes)

PRIORITY	COLOR	CONDITION	NOTES
1	Red	Immediate	Life threatening
2	Yellow	Urgent	Can delay up to 1 hr.
3	Green	Delayed	Up to 3 hours
4	Black	Deceased	No care needed

#### Priority 1 - Immediate transport:

Unconscious, disoriented, very confused, rapid respirations, weak irregular pulse, severe uncontrolled bleeding, other signs of shock (cold, clammy skin, low blood pressure, etc.).

#### Priority 2 - Urgent, can delay transport up to 1 hour:

Conscious, oriented, with any significant fracture or other significant injury, but without signs of shock.

#### Priority 3 - Delayed transport up to 3 hours:

Walking wounded, CAO x3, minor injuries.

#### Priority 4 - Deceased, no care needed:

No pulse, no respirations (open airway first, obvious mortal wounds (e.g. decapitation).

#### NOTES

**Assessment of patients should be <1 minute\* each.**

\*(Have someone else control bleeding during your survey.)

**All unconscious patients are Priority 1 - Immediate.**

"Walking wounded" are usually GREEN - Priority 3.

**All pulseless patients are BLACK - Priority 4.**